Revised November 2013



Table 1. Treatment Response of Common Aquatic Plants to Registered Herbicides

	bispyribac	carfentrazone	copper & copper complexes -	copper complexes herbicides	diquat	endothall	fluridone	flumioxazin	glyphosate	imazamox	imazapyr	penoxsulam	sodium carbonate	peroxy-nyarate triclopyr	2,4-D	
Aquatic Group							Aquati	c Herk	oicide¹							Grass Carp ⁹
& vegetation				- 25			-				E					
						- 2 2		gae								
Chara/Nitella	Р		E		Р	G ² -P ³	Р	P	P		- 1		1		P	G
filamentous			E		G	G ² -P ³	Р	G	P				G ⁶		Р	F
planktonic			E		Р	G²	Р	F	Р		- 0		G ⁶		Р	
			1	- 33			Floatin	g Plar	nts				<i>[</i>			
azolla		G	P		G		E	E	F			E			F	
duckweeds		E	Р		G	Р	E	E	P		Р	E/I			F	F
salvinia	F	G	Р		G		E	E	G	E		E				Р
water hyacinth	E	G	Р	G ⁴	ΛE		Р	P	G	E	EV	E		E	E	Р
watermeal	F	G	Р		F		G	E				G			F	Р
water lettuce	E	// E	P	G ⁴	E		G	E	G		E	E		G	F	
			L				Submer	ged pla	ants		-	1		195		_
coontail	Р		Р	G ⁴	Е	E	E	G				V			G	F- G
elodea			P	G ⁴	E	F	E	E				G				E
fanwort			Р	Р	G	F	E	G				G			F	F
hydrilla	E		Р	G⁴	G	G	E	G		G		Ε				E
milfoils	G	E	Р	G⁴	E	E	G	G		G		Ε		Ε	E	F
naiads			Р	G⁴	E	E	Ε	E,d				G			F	E
parrotfeather			Р	Р	E	E	E	G		G	G⁵	G		G	E	G
pondweeds	G		Р	G ⁴	G	E	E	G		E	G⁵	G			Р	E

¹E= excellent control; G= good control; F= fair control; P= poor control; blank= unknown or no control

²Hydrothol formulations

⁶Best on blue-green algae

³Aquathol formulations

⁷E for sedge

⁴Specific copper complexes only- Nautique, Komeen, etc.

⁸F for rushes

⁵Spray only emergent portion

⁹Permit required from Texas Parks & Wildlife

TEXAS A&M
GRILIFE
EXTENSION

	ribac ntrazone		<u> </u>	5	thall	thail	oxazin	osate	атох	apyr	penoxsulam	um carbonate	cy-hydrate	pyr		KILIF	
	bispy	carfe	coppe	copp - her	diquat	endo	flurido	flumio	glyph	imaza	imazapy	peno	sodiu	pero	triclopy	2,4-D	
quatic Group & vegetation				1.5			Aquati	ic Herbic	ides¹								Grass Ca
							Emorgo	nt Dlant	_								

Aquatic Group & vegetation							Aquatio	Her	rbici	des ¹							Grass Carp ⁹
							Emergen	t Pla	ants	,							
alders			Р		F	Р	Р			E		E			E	E	
alligatorweed	E	F			Р		F	G		G	G	E			E	F	
arrowhead	E		Р		G	G	E	G		E	E	E				E	
buttonbrush			Р		F	Р	Р			G		G				F	
cattails	Р		Р		G	Р	F	Р		E	E	E				F	
common reed			P		F		F	Р		E	G	E	\ /			F	
frogbit	E		<i>[</i>	F ⁴	E			G		F	E	E			E	E	
pickerelweed	F			F ⁴	G		Р	Р		F	E	E			G	G	
sedges & rushes	F		P		F		Р	F		G		E ⁷ F ⁸	G			F	
slender spikerush			Р		G		G	Р		Р		F					
smartweed	G		Р	F ⁴	F		F	Р		E	E	E	G		E	E	
southern watergrass			Р				G			E		E				Р	
waterlilies	F		P		Р		E	F		G	G	G	G		G	E	
water pennywort	G		P		G		Р	G		G		E	G		E	G	
water primrose		F	P		F	P	F	G		E	E	E			E	E	
watershield			P		Р		G	G		G	G	E				E	
willows	Р		Р		F	Р	Р	Р		E		Е			E	E	
Active Ingredients	Commonly Available Trade Names								Ac	tive Ing	<u>re</u> dient	:s	Common	ly Ava	ilable Tra	de Namo	<u>es</u>

bispyribac	Tradewind		glyphosate	Rodeo, Aquamaster, AquaNeat, Eraser AQ, Refuge®, others				
carfentrazone	Stingray		imazamox	Clearcast				
copper &	Copper Sulfate, Cutrine, Cutrine Plus, K-Te	ea, Captain, Agritec,	imazapyr	Habitat, Arsenal, Poloris				
complexes	EarthTec, Clearigate		penoxsulam	Galleon				
copper - herbicides	Komeen, Nautique		sodium carbonate	Green Clean, PAK 27, Phycomycin				
diquat	Reward, Harvester, Tribune, Tsunami DQ,	Diquat SPC2L, Weedtri	in φ Đroxyhydrate	Green Clean, PAK 27, Phycomychi				
endothall	Aquathol K, Aquathol Super K, Hydrothol	191	triclopyr	Renovate, Navitrol, Ecotriclopyr				
flumioxazin	Clipper		2,4-D	Navigate, Weedar 64				
fluridone	Sonar, Avast, WhiteCap, Restore		***Tovas A&M Univer	sity & Extension does not endorse any trade name herbicide				

^{***}Texas A&M University & Extension does not endorse any trade name herbicide



Table 2. Aquatic Vegetation Herbicide Control Water Use Restriction¹ (number of days after treatment before use in private waters only)

		Human Use		Livestock		Irrigation		
Common Name	Drinking	Swimming	Fish	Watering	-21	Turf	Crops	
bispyribac	0	0	0	0		30	30	
carfentrazone	0 - 1 ²	0	0	0 - 1 ²		0 - 14 ²	0 - 14²	
copper complexes ³	0	0	0	0		0	0	
diquat	1-3 ³	0	0	1		1-34	5	
endothall ⁵	7-25	1	0	7-25		7-25	7-25	
flumioxazin	0	0	0	0		0-34	5	
fluridone ⁶	0	0	0	0		7-30	7-30	
glyphosate ⁷	0	0	0	0		0	0	
imazamox	0	0	0	0		1	1 ⁸	
imazapyr	*9	0	0	0		120 ¹⁰	120 ¹⁰	
penoxsulam	0	0	0	0		0	*11	
SCP ¹²	0	0	0	0	- //	0	0	
triclopyr	*13	0	0	0		014	120 ¹⁵	
2,4-D	*16	*16	*16	*16		*16	*16	

¹ Aquatic vegetation control can result in period of low dissolved oxygen which can stress and/or kill fish. It is best to treat most aquatic vegetation early in the growing season, when the plant is rapidly growing. Treating small areas (e.g. 1/4) of pond at a time at 10-14 day intervals will allow for decomposition usually without causing oxygen depletion.

ONLY PRODUCTS LABELED FOR AQUATIC USE may be used in, over, or near the water

Additional information is available through the following references and websites – aquaplant.tamu.edu, srac.tamu.edu, & wildlife.tamu.edu

Aquatic Vegetation Indentification Card Deck - Pub. #B6095, produced by Dr. Michael P. Masser are for sale for \$12.00 + taxes & shipping, order for 10 or more or \$7.00+ Plus taxes & shipping, order from the Texas AgriLife Bookstore,

These tables were prepared and maintained by:
Michael P. Masser, Professor and Department Head WFSC
Todd D. Sink, Assistant Professor and Fisheries Extension Specialist

² Varies if 20% or more of surface area is treated

³ If water is for drinking, the elemental copper concentration should not exceed 1.0 ppm (i.e. 4.0 ppm copper sulfate).

⁴ Depending on formulation or rate - **Read label**.

⁵ Length of use restriction for endothall varies with concentration used. **Read label**.

⁶ Do not apply within 0.25 mile of a functioning potable water intake.

⁷ Do not apply within 0.5 mile of a functioning potable water intake.

⁸ Do not use treated water to irrigate greenhouses, nurseries, or hydroponics

⁹ Greater than 1/2 mile from potable water intake

¹⁰ Or until <1.0 ppb

¹¹Do not use water from any treated site for food crop irrigation until residues are determined to be less than or equal to 1 ppb.

¹² Sodium Carbonate Peroxyhydrate

¹³ Minimum setback distances from potable water intakes required and laboratory tests to determine < 0.4 ppm for use. **Read label**.

¹⁴ No restriction on irrigating established grasses but do not harvest hay for 14 days after application. **Read label**.

¹⁵ Or until non-detectable concentration in immunoassay analysis

¹⁶ Water restrictions on 2,4-D vary with formulation, location, rate, and time of year. **Read label.**

<agrilifebookstore.org> or fax 979/458-0172

^{*}srac.tamu.edu website publication numbers, SRAC 0360-0369; 3600-3699